



First record of the genus *Syritta* Le Peletier & Audinet-Serville, 1828 (Diptera, Syrphidae) from the West Indies

Ximo Mengual¹, Carlos de Soto Molinari²

¹ Zoologisches Forschungsmuseum Alexander Koenig, Leibniz-Institut für Biodiversität der Tiere, Adenauerallee 160, Bonn, D-53113, Germany.

² Calle K, No. 2, esquina 13, Altos Arroyo Hondo, Santo Domingo, Dominican Republic.

Corresponding author: Ximo Mengual, x.mengual@leibniz-zfmk.de

Abstract

We report the genus *Syritta* Le Peletier & Audinet-Serville, 1828 (Diptera, Syrphidae) from West Indies for the first time. A female of the synanthropic species *Syritta flaviventris* Macquart, 1842 was photographed in the Refugio de Vida Silvestre Río Higuamo, Dominican Republic. This species was introduced in the American continent by human expansion.

Keywords

Dominican Republic, flower fly, hover fly, photographic record, social media, synanthropic.

Academic editor: Alexandre Pereira-Colavite | Received 8 June 2020 | Accepted 23 July 2020 | Published 10 August 2020

Citation: Mengual X, de Soto Molinari C (2020) First record of the genus *Syritta* Le Peletier & Audinet-Serville, 1828 (Diptera, Syrphidae) from the West Indies. Check List 16 (4): 991–995. <https://doi.org/10.15560/16.4.991>

Introduction

Commonly known as flower flies or hover flies, members of the family Syrphidae (Insecta, Diptera) are diverse in number (more than 6,200 described species) and larval biology (Rotheray and Gilbert 2011). These conspicuous flies provide important ecosystems services since adults are crucial pollinators of crops and wild plants (Inouye et al. 2015; Rader et al. 2016) and larvae play important roles as biological control agents (Tenhumberg and Poehling 1995; Rojo et al. 2003; Grosskopf 2005) and organic matter recyclers (Lardé 1989; Pérez-Bañón et al. 2013).

The flower fly genus *Syritta* Le Peletier & Audinet-Serville, 1828 (Diptera, Syrphidae, Eristalinae) comprises 61 described species occurring in all biogeographical regions, except both poles, and its highest number of species is found in the Afrotropical Region, with 43 (Lyneborg and Barkemeyer 2005; Van Steenis 2010).

They are small-sized flower flies, usually with a black and yellow-orange to brown coloration, with enlarged hind femora. While adults visit flowers to feed on pollen and nectar (Gilbert 1981), larvae are saprophagous and feed on decaying plant and animal matter (Pérez-Bañón and Marcos-García 2000; Magni et al. 2013).

Two *Syritta* species are present in the Nearctic and the Neotropical Regions, i.e., *S. pipiens* (Linnaeus, 1758) and *S. flaviventris* Macquart, 1842, both introduced as a result of human expansion (Thompson et al. 1990; Lyneborg and Barkemeyer 2005; Skevington et al. 2019). These two synanthropic species can be easily distinguished based on the morphological characteristics provided by Thompson et al. (1990) and Skevington et al. (2019). Pérez-Bañón and Marcos-García (2000) described the larvae of *S. flaviventris* from *Opuntia* Mill.

platyclades and compared them with the larvae of *S. pipiens*. Records of both species are abundant in online databases, such as BugGuide (www.bugguide.net) and iNaturalist (www.inaturalist.org), and field observations in the New World are graphically documented in these databases and incorporated in the Global Biodiversity Information Facility (www.gbif.org).

Despite the available information and different ways of information dissemination, the genus *Syritta* has not been yet recorded from the West Indies (Thompson 1981; Perez-Gelabert 2008, 2020). Here we report the first photographic record of *S. flaviventris* from a Caribbean island, Hispaniola.

Methods

A photograph of a female flower fly (Fig. 1) was taken on 17 February 2013 by the second author in the Refugio de Vida Silvestre Río Higuamo, in the San Pedro Macorís province, located in the south-east of the Dominican Republic around the Higuamo river. The area is part of the Hispaniolan moist forest, a tropical and subtropical moist broadleaf forest ecoregion on the island of Hispaniola (Ecoregions 2017). The photograph was posted on a Facebook group (Insectos Caribeños 2020) and rapidly got the attention of the first author.

SimpleMappr (Shorthouse 2010) was used to create the map. Records (point data) were obtained from GBIF (2020) and published literature (Fluke 1960;

Marnef 1967; Campos and Peña 1973; Thompson et al. 1976; Thompson et al. 1990; Morales and Köhler 2008; Ramírez-García 2010; Eloisa 2014; Sidhu and Biddinger 2014; Flores 2017; Skevington et al. 2019; Montes 2020).

Results

The photographed female was identified as *Syritta flaviventris* Macquart, 1842 (Fig. 1), representing the first record of the genus *Syritta* for the West Indies.

New records. DOMINICAN REPUBLIC • 1 ♀; San Pedro de Macorís, Refugio de Vida Silvestre Río Higuamo; 18.47169°N, 069.32667°W; 4 m a.s.l.; 17 Feb. 2013; photographed by C de Soto Molinari.

Identification. The most obvious diagnostic characteristic of *S. flaviventris* is the wing without spurious vein, but the wing venation is not in focus in the photograph (Fig. 1). Another good characteristic that distinguish *S. flaviventris* from *S. pipiens* is the hind femur with a basoventral tubercle (large in males, smaller in females; absent in *S. pipiens*), but it is also not visible because the picture was taken in dorsal view. Our identification was based on the following morphological characteristics stated by Thompson et al. (1990): face silvery white pruinose, antenna extensively dark, pruinose markings near transverse suture on scutum, fore and mid legs entirely orange, and wing veins orange.



Figure 1. First record of the genus *Syritta* from West Indies. Female of *Syritta flaviventris* photographed in Refugio de Vida Silvestre Río Higuamo, the Dominican Republic.




Figure 2. Records of *Syritta flaviventris* in the American continent. Blue circles are records from GBIF and published literature; the red circle is our new record from the Dominican Republic. **A.** Overview of the American continent with the records of *S. flaviventris*; the grey area is enlarged below. **B.** Records in the Caribbean region, between longitudes of 011.5°N and 030.5°N.

Discussion

In the New World, *Syritta flaviventris* has been reported from several states in the United States (Thompson et al. 1990; Sidhu and Biddinger 2014; Skevington et al. 2019; GBIF 2020), Mexico (Colima: Ramírez-García 2010; GBIF 2020; Mexico State: Eloisa 2014; Flores 2017; and Nuevo León: Thompson et al. 1990; GBIF 2020), Brazil (São Paulo: Fluke 1960; Rio Grande do Sul: Morales and Köhler 2008), Uruguay (Montevideo: GBIF 2020), Argentina (Buenos Aires, Santa Fé and Entre Ríos: Thompson et al. 1976; GBIF 2020), and Chile (Valparaíso and Metropolitan region: Marnef 1967, as *Austrosyritta cortesi* Marnef, 1967, junior synonym of *S. flaviventris*; and Biobío: Montes 2020). Campos and Peña (1973) recorded *S. flaviventris* from Easter Island.

Our photographic record represents the first report from the Dominican Republic of this introduced genus (Perez-Gelabert 2008, 2020) and from the whole West Indies (Thompson 1981). Our record is 1,400 km away from the closest known record for this species in Miami-Dade County, Florida, United States (Lisnel 2019) (Fig. 2). Moreover, it is the first record of this genus on an island in the Americas, besides Easter Island in the Pacific Ocean, and points out the survival and establishment capacities of this species in new environments.

This is not the first flower fly report based on a photographic record. Álvarez Fidalgo et al. (2018) reported *Leucozona laternaria* (Müller, 1776) from the Iberian Peninsula based on several photographic records. Online databases and networks, like iNaturalist and BugGuide, images repositories such as Flickr (www.flickr.com), and

social media like Facebook (www.facebook.com), offer new ways of information dissemination and high-quality geographical data for taxonomists. These digital tools can provide new insights on distributional data, first graphical documents of rare species or even be used to describe new species based on images posted in these websites and social media (Winterton et al. 2012; Amézquita et al. 2013; Gonella et al. 2015; Silverman 2016; Winterton 2020). Social media and online networks are growing in importance for citizen science projects (Tiago et al. 2017; Irwin 2018) and biologists should engage, collaborate and benefit from the public with scientific concerns and willing to enact (Chandler et al. 2017; Suprayitno et al. 2017; von Konrat et al. 2018).

Acknowledgements

We want to thank Daniel E. Perez-Gelabert for creating the Facebook group “Insectos Caribeños” and for promoting the study of insects in the Caribbean. We thank Augusto L. Montoya, Gil F. Miranda and Martin Hauser for their helpful comments and suggestions.

Authors' Contributions

XM identified the specimen and wrote the manuscript, and CdSM photographed the specimen and commented on the manuscript.

References

- Álvarez Fidalgo M, Álvarez Fidalgo P, Ricarte A, Marcos-García MÁ (2018) The genus *Leucozona* Schiner, 1860 on the Iberian Peninsula, including the first records of *Leucozona laternaria* (Müller, 1776) (Diptera: Syrphidae). *BV news Publicaciones Científicas* 7 (98): 128–141.
- Amézquita A, Márquez R, Medina R, Mejía-Vargas D, Kahn TR, Suárez G, Mazariegos L (2013) A new species of Andean poison frog, *Andinobates* (Anura: Dendrobatidae), from the northwestern Andes of Colombia. *Zootaxa* 3620 (1): 163–178. <https://doi.org/10.11646/zootaxa.3620.1.8>
- Campos L, Peña LE (1973) Los insectos de Isla de Pascua (Resultados de una prospección entomológica). *Revista Chilena de Entomología* 7: 217–229.
- Chandler M, See L, Copas K, Bonde AMZ, Claramunt López B, Danielsen F, Legind JK, Masinde S, Miller-Rushing AJ, Newman G, Rosemartin A, Turak E (2017) Contribution of citizen science towards international biodiversity monitoring. *Biological Conservation* 213 (Part B): 280–294. <https://doi.org/10.1016/j.biocon.2016.09.004>
- Ecoregions (2017) <https://ecoregions2017.appspot.com/>. Accessed on: 2020-7-21.
- Eloisa A (2014) *Syritta flaviventris*. <https://www.inaturalist.org/observations/1084778>. Accessed on: 2020-6-26.
- Flores E (2017) *Syritta flaviventris*. <https://www.inaturalist.org/observations/5670221>. Accessed on: 2020-6-26.
- Fluke CL (1960) Concerning the Catalogue of Neotropical Syrphidae. *Revista Brasileira de Entomologia* 9: 169.
- GBIF (2020) GBIF Occurrence Download. <https://doi.org/10.15468/dl.6dee2h>. Accessed on: 2020-6-26.
- Gilbert FS (1981) Foraging ecology of hoverflies: morphology of the mouthparts in relation to feeding on nectar and pollen in some common urban species. *Ecological Entomology* 6 (3): 245–262. <https://doi.org/10.1111/j.1365-2311.1981.tb00612.x>
- Gonella PM, Rivadavia F, Fleischmann A (2015) *Drosera magnifica* (Droseraceae): the largest New World sundew, discovered on Facebook. *Phytotaxa* 220 (3): 257–267. <https://doi.org/10.11646/phytotaxa.220.3.4>
- Grosskopf G (2005) Biology and life history of *Cheilosia urbana* (Meigen) and *Cheilosia psilophthalma* (Becker), two sympatric hoverflies approved for the biological control of hawkweeds (*Hieracium* spp.) in New Zealand. *Biological Control* 35 (2): 142–154. <https://doi.org/10.1016/j.biocontrol.2005.06.013>
- Inouye DW, Larson BMH, Ssymank A, Kevan PG (2015) Flies and flowers III: Ecology of foraging and pollination. *Journal of Pollination Ecology* 16 (16): 115–133. [http://doi.org/10.26786/1920-7603\(2015\)15](http://doi.org/10.26786/1920-7603(2015)15)
- Insectos Caribeños (2020) <https://www.facebook.com/groups/1461724150742197/>. Accessed on: 2020-6-26.
- Irwin A (2018) Citizen science comes of age. *Nature* 562: 480–482. <https://doi.org/10.1038/d41586-018-07106-5>
- Lardé G (1989) Investigation on some factors affecting larval growth in a coffee-pulp bed. *Biological Wastes* 30 (1): 11–19. [https://doi.org/10.1016/0269-7483\(89\)90139-0](https://doi.org/10.1016/0269-7483(89)90139-0)
- Le Peletier ALM, Audinet-Serville JG (1828) *Xylote*, *Xylota*. In: Latreille PA (Ed) *Encyclopédie Méthodique, Histoire Naturelle. Entomologie, ou histoire naturelle des Crustacés, des Arachnides et des Insectes*, Tome 10. Agasse, Paris, 808–809.
- Linnaeus C (1758) *Systema Naturae per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis* (10th ed). Laurentii Salvii, Holmiae [= Stockholm], 1384 pp. <https://doi.org/10.5962/bhl.title.542>
- Lisnel (2019) *Syritta flaviventris*. <https://www.inaturalist.org/observations/27099564>. Accessed on: 2020-6-26.
- Lyneborg L, Barkemeyer W (2005) The Genus *Syritta*, A World Revision of the Genus *Syritta* Le Pelletier & Serville, 1828 (Diptera: Syrphidae). *Entomograph* 15. Apollo Books, Denmark, 224 pp.
- Macquart PJM (1842) *Diptères exotiques nouveaux ou peu connus*. Tome deuxième, 2^e partie. Mémoires de la Société royale des Sciences, de l'Agriculture et des Arts de Lille 1841 (1): 65–200, 22 pls.
- Magni PA, Pérez-Bañón C, Borrini M, Dadour IR (2013) *Syritta pipiens* (Diptera: Syrphidae), a new species associated with human cadavers. *Forensic Science International* 231 (1–3): e19–e23. <https://doi.org/10.1016/j.forsciint.2013.05.023>
- Marnef L (1967) *Austrosyritta cortesi* nov. gen., nov. sp. de sirfido de Chile (Diptera Syrphidae). *Bulletin et Annales de la Société Royale Belge d'Entomologie* 103: 268–275.
- Montes O (2020) *Syritta flaviventris*. <https://www.inaturalist.org/observations/37455564>. Accessed on: 2020-6-26.
- Morales MN, Köhler A (2008) Comunidade de Syrphidae (Diptera): diversidade e preferências florais no Cinturão Verde (Santa Cruz do Sul, RS, Brasil). *Revista Brasileira de Entomologia* 52 (1): 41–49. <https://doi.org/10.1590/S0085-56262008000100008>
- Pérez-Bañón C, Hurtado P, García-Gras E, Rojo S (2013) SEM studies on immature stages of the drone flies (Diptera, Syrphidae): *Eristalis similis* (Fallen, 1817) and *Eristalis tenax* (Linnaeus, 1758). *Microscopy Research and Technique* 76 (8): 853–861. <https://doi.org/10.1002/jemt.22239>
- Pérez-Bañón C, Marcos-García MA (2000) Description of the immature stages of *Syritta flaviventris* (Diptera: Syrphidae) and new data about the life history of European species of *Syritta* on *Opuntia maxima*. *European Journal of Entomology* 97 (1): 131–136. <https://doi.org/10.14411/eje.2000.022>
- Perez-Gelabert DE (2008) Arthropods of Hispaniola (Dominican Republic and Haiti): a checklist and bibliography. *Zootaxa* 1831 (1): 1–530. <https://doi.org/10.11646/zootaxa.1831.1.1>
- Perez-Gelabert DE (2020) Checklist, bibliography and quantitative data of the arthropods of Hispaniola. *Zootaxa* 4749 (1): 1–668. <https://doi.org/10.11646/zootaxa.4749.1.1>

- Rader R, Bartomeus I, Garibaldi LA, Garratt MP, Howlett BG, Winfree R, Cunningham SA, Mayfield MM, Arthur AD, Andersson GK, Bommarco R, Brittain C, Carvalheiro LG, Chacoff NP, Entling MH, Foully B, Freitas BM, Gemmill-Herren B, Ghazoul J, Griffin SR, Gross CL, Herbertsson L, Herzog F, Hipólito J, Jagger S, Jauker F, Klein AM, Kleijn D, Krishnan S, Lemos CQ, Lindström SA, Mandelik Y, Monteiro VM, Nelson W, Nilsson L, Pattemore DE, Pereira Nde O, Pisanty G, Potts SG, Reemer M, Rundlöf M, Sheffield CS, Scheper J, Schüepp C, Smith HG, Stanley DA, Stout JC, Szentgyörgyi H, Taki H, Vergara CH, Viana BF, Woyciechowski M (2016) Non-bee insects are important contributors to global crop pollination. *Proceedings of the National Academy of Sciences of the United States of America* 113 (1): 146–151. <https://doi.org/10.1073/pnas.1517092112>
- Ramírez-García E (2010) *Syritta flaviventris* Macquart, 1842. <http://unibio.unam.mx/irekani/handle/123456789/26112?proyecto=Irekani>. Accessed on: 2020-6-26.
- Rojo S, Gilbert FS, Marcos-García MA, Nieto JM, Mier MP (2003) A World Review of Predatory Hoverflies (Diptera, Syrphidae: Syrphinae) and Their Prey. *Centro Iberoamericano de la Biodiversidad*, Alicante, 319 pp.
- Rotheray GE, Gilbert F (2011) The natural history of hoverflies. *Forest Text*, Ceredigion, 333 pp.
- Shorthouse DP (2010) SimpleMappr, an online tool to produce publication-quality point maps <https://www.simplemappr.net>. Accessed on: 2020-6-26.
- Sidhu CS, Biddinger DJ (2014) Northern-most North American flower visitation records of the introduced flower fly, *Syritta flaviventris* (Diptera: Syrphidae) and comparisons with sympatric species, *Syritta pipiens*. *The Great Lakes Entomologist* 47 (3–4): 166–170.
- Silverman L (2016) <https://www.npr.org/sections/alltechconsidered/2016/08/06/488830352/the-app-that-aims-to-gamify-biology-has-amateurs-discovering-new-species>. Accessed on: 2020-5-8.
- Skevington J, Locke M, Young A, Moran K, Crins W, Marshall S (2019) Field guide to the flower flies of northeastern North America. Princeton University Press, Princeton, 512 pp.
- Suprayitno N, Narakusumo R, von Rintelen T, Hendrich L, Balke M (2017) Taxonomy and Biogeography without frontiers – WhatsApp, Facebook and smartphone digital photography let citizen scientists in more remote localities step out of the dark. *Biodiversity Data Journal* 5: e19938. <https://doi.org/10.3897/BDJ.5.e19938>
- Tenhumberg B, Poehling H-M (1995) Syrphids as natural enemies of cereal aphids in Germany: aspects of their biology and efficacy in different years and regions. *Agriculture, Ecosystems and Environment* 52 (1): 39–43. [https://doi.org/10.1016/0167-8809\(94\)09007-t](https://doi.org/10.1016/0167-8809(94)09007-t)
- Thompson FC (1981) The flower flies of the West Indies (Diptera: Syrphidae). *Memoirs of the Entomological Society of Washington* 9: 1–200.
- Thompson FC, Fee FD, Berzark LG (1990) Two immigrant synanthropic flower flies (Diptera: Syrphidae) new to North America. *Entomological News* 101 (2): 69–74.
- Thompson FC, Vockeroth JR, Sedman YS (1976) 46. Family Syrphidae. A catalogue of the Diptera of the Americas south of the United States. *Museu de Zoologia, Universidade de São Paulo, São Paulo*, 195 pp.
- Tiago P, Ceia-Hasse A, Marques TA, Capinha C, Pereira HM (2017) Spatial distribution of citizen science casuistic observations for different taxonomic groups. *Scientific Reports* 7: 12832. <https://doi.org/10.1038/s41598-017-13130-8>
- Van Steenis J (2010) A new species of the genus *Syritta* Le Peletier & Serville, 1828 (Diptera, Syrphidae), with new distributional records of other *Syritta* species. *Norwegian Journal of Entomology* 57: 111–119.
- von Konrat M, Campbell T, Carter B, Greif M, Bryson M, Larrain J, Trouille L, Cohen S, Gaus E, Qazi A, Ribbens E, Livshultz T, Walker TJ, Suwa T, Peterson T, Rodriguez Y, Vaughn C, Yang C, Aburahmeh S, Carstensen B, de Lange P, Delavoi C, Strauss K, Drag J, Aguero B, Snyder C, Martinec J, Smith A (2018) Using citizen science to bridge taxonomic discovery with education and outreach. *Applications in Plant Sciences* 6 (2): e1023. <https://doi.org/10.1002/aps3.1023>
- Winterton S (2020) A new bee-mimicking stiletto fly (Therevidae) from China discovered on iNaturalist. *Zootaxa* 4816 (3): 361–369. <https://doi.org/10.11646/zootaxa.4816.3.6>
- Winterton S, Guek H, Brooks S (2012) A charismatic new species of green lacewing discovered in Malaysia (Neuroptera, Chrysopidae): the confluence of citizen scientist, online image database and cybertaxonomy. *ZooKeys* 214: 1–11. <https://doi.org/10.3897/zookeys.214.3220>